

Environmental Recommendations

0 - 20 weeks of age

Floor (sq. ft.)	1.0	0 - 8 weeks of age
	2.0	8 - 20 weeks of age
Feeder (in.)	3.0	or 20 - 25 birds per pen
Water (in.)	1.0	or 50 - 100 birds per round metal drinker

20 + weeks of age

Floor (sq. ft.)	1.5	White egg breeds
	2.0	Brown egg breeds
Feeder (in.)	3.0	or 20 birds per pen
Water (in.)	2.0	or 50 - 100 birds per round metal drinker

Broilers

Floor (sq. ft.)	1.0 - 2.5	Increases if birds are fed beyond 8 weeks
Feeder (in.)	3.0	or 20 - 25 birds per pen
Water (in.)	1.0	or 50 - 100 birds per round metal drinker



Eldora • Dyersville
www.unitedsuppliers.com

Farm Flock

A complete feeding program for starting pullets, layers, & broilers.



20% Broiler Starter

Complete ration designed to meet the demands of young, growing chicks. Medicated to increase feed efficiency & aid in growth promotion.

Bacitracin Methylene Disalicylate...	30 g/ton
Amprolium.....	0.0125%
Protein, min	20%
Lysine, min	1.25%
Methionine, min	0.52%
Fat, min	5%
Fiber, max	4.5%
Calcium.....	1%
Phosphorus, min	0.78%
Vitamin A, min	4,500 IU/lb
Vitamin D, min.....	2,000 IU/lb
Vitamin E, min.....	15 IU/lb
Vitamin K, min	1.0 mg/lb
Vitamin C, min	91 mg/lb

Directions for use: Feed as sole ration to chicks from hatching to 5-6 weeks of age. Each chick will consume 8-10 pounds of Broiler Starter. Switch to 18% Broiler Finisher or Broiler Concentrate thereafter.



18% Broiler Finisher

Convenient feed designed to be fed from 5-6 weeks until slaughter.

Protein, min	18%
Lysine, min	1.06%
Methionine, min	0.47%
Fat, min	5%
Fiber, max	4%
Calcium.....	0.90%
Phosphorus, min	0.85%
Vitamin A, min	4,500 IU/lb
Vitamin D, min.....	1,900 IU/lb
Vitamin E, min.....	10 IU/lb
Vitamin K, min	1.0 mg/lb
Vitamin C, min	90 mg/lb

Directions for use: Feed as sole ration to broiler chicks 5-6 weeks of age through slaughter. Each broiler chick will consumer 10-12 pounds of Broiler Finisher

Broiler Concentrate Medicated

Grind & mix concentrate designed to meet the demands of young, growing broilers.

Bacitracin Methylene Disalicylate...	90 g/ton
Amprolium.....	0.0358%
Protein, min	37%
Lysine, min	2.45%
Methionine, min	0.93%
Fat, min	8%
Fiber, max	4.5%
Calcium.....	2.3%
Phosphorus, min	1.35%
Vitamin A, min	10,000 IU/lb
Vitamin D, min.....	5,000 IU/lb
Vitamin E, min.....	35 IU/lb
Vitamin K, min	2.4 mg/lb
Vitamin C, min	225 mg/lb

Directions for use: Use following grind & mix instructions to achieve starter and grower rations.

	Starter/Grower	Grower/Finisher
Cracked Corn	1,100 lbs	1,300 lbs
Broiler Concentrate	900 lbs	700 lbs
	2,000 lbs	2,000 lbs

26% Free Choice Balancer

Complete balancer formulated to compliment available feed-stuffs such as corn, oats, grit & oyster shells.

Protein, min	26%
Lysine, min	1.5%
Methionine, min	0.42%
Calcium, min	3.25%
Calcium, max.....	3.75%
Phosphorus, min	1.60%

Directions for use: Feed along with corn and oats on a free choice basis. Grit & oyster shells may be offered free choice if desired.

Complete Layer 17-E

Complete layer ration designed to meet the needs of egg producing layers.

Protein, min	16%
Lysine, min	0.75%
Methionine, min	0.30%
Calcium, min	3.5%
Calcium, max.....	4%
Phosphorus, min	0.65%

Directions for use: Feed as the only grain source free choice to laying hens. Layers will consume 20 - 23 lbs per 100 birds per day.

Layer Concentrate 40-E

Complete layer ration designed to meet the needs of egg producing layers.

Protein, min	36%
Lysine, min	2.40%
Methionine, min	0.80%
Calcium, min	4%
Calcium, max.....	4.50%
Phosphorus, min	2%

	Grower 6-14 weeks	Developer 14 weeks to pre-lay	Phase 1	Phase 2	Phase 3
Cracked Corn, lbs	1375	1325	1375	1400	1400
Ground Oats, lbs	100	200	---	---	---
40-E Concentrate, lbs	525	475	500	450	425
Limestone/Oyster Shells, lbs	---	---	125	150	175
Equivalent Crude Protein	16.5%	16%	17%	16%	15%
Calcium	---	---	3.2%	3.3%	3.8%

Directions for use: Feed as the sole ration to growing & developing pullets as well as laying hens. Laying hens will consume 20 - 23 pounds of total ration per 100 birds per day.